

OBSTETRIC HEMORRHAGE

# Risk Assessment Tables

## LABOR & DELIVERY ADMISSION

	MEDIUM RISK	HIGH RISK
<b>RISK FACTORS</b>	<input type="checkbox"/> Prior cesarean, uterine surgery, or multiple laparotomies	<input type="checkbox"/> Placenta previa/low lying
	<input type="checkbox"/> Multiple gestation	<input type="checkbox"/> Suspected accreta/percreta
	<input type="checkbox"/> > 4 prior births	<input type="checkbox"/> Platelet count < 70,000
	<input type="checkbox"/> Prior PPH	<input type="checkbox"/> Active bleeding
	<input type="checkbox"/> Large myomas	<input type="checkbox"/> Known coagulopathy
	<input type="checkbox"/> EFW > 4000 g	<input type="checkbox"/> 2 or more medium risk factors
	<input type="checkbox"/> Obesity (BMI > 40)	/
	<input type="checkbox"/> Hematocrit < 30% & other risk	/
<b>INTERVENTION</b>	<input type="checkbox"/> Type & SCREEN, review protocol	<input type="checkbox"/> Type & CROSS, review protocol

## INTRAPARTUM

	MEDIUM RISK	HIGH RISK
<b>RISK FACTORS</b>	<input type="checkbox"/> Chorioamnionitis	<input type="checkbox"/> New active bleeding
	<input type="checkbox"/> Prolonged oxytocin > 24 hours	<input type="checkbox"/> 2 or more medium (admission and/or intrapartum) risk factors
	<input type="checkbox"/> Prolonged 2nd stage	/
	<input type="checkbox"/> Magnesium sulfate	/
<b>INTERVENTION</b>	<input type="checkbox"/> Type & SCREEN, review protocol	<input type="checkbox"/> Type & CROSS, review protocol

\* Establish a culture of huddles for high-risk patients and post-event debriefing \*